

Santos VS, Gurgel RQ, Cuevas LE, Martins-Filho PR. Prolonged fecal shedding of SARS-CoV-2 in pediatric patients. A quantitative evidence synthesis. *J Pediatr Gastroenterol*

Table 3 Santos 2020. Included references

<p>Xu Y, Li X, Zhu B, et al. Characteristics of pediatric SARS-CoV-2 infection and potential evidence for persistent fecal viral shedding. <i>Nat Med</i> 2020; 26 (4):502–505. doi:10.1038/s41591-020-0817-4.</p>
<p>Cai J, Xu J, Lin D, et al. A Case Series of children with 2019 novel coronavirus infection: clinical and epidemiological features. <i>Clin Infect Dis</i> 2020; doi:10.1093/cid/ciaa198.</p>
<p>Ma X, Su L, Zhang Y, Zhang X, Gai Z, Zhang Z. Do children need a longer time to shed SARS-CoV-2 in stool than adults? <i>J Microbiol Immunol Infect</i> 2020; doi:10.1016/j.jmii.2020.03.010.</p>
<p>Tan YP, Tan BY, Pan J, Wu J, Zeng SZ, Wei HY. Epidemiologic and clinical characteristics of 10 children with coronavirus disease 2019 in Changsha, China. <i>J Clin Virol</i> 2020; 127:104353.doi:10.1016/j.jcv.2020.104353.</p>